

Document Title: Description, AN, How LP-80 measures LAI		Part # and Rev. 13381	
		Release Date: 1-12-07	
Rev.	Description	Revision By	Date

Production Filename: 13381 (In Product Library)

Path to Working Files: DecaDoc\Application Notes\Master\Description

Dimensions: 8.5 inch wide, 11 inch tall

Material: Paper, 92 Bright White or better, 75g/m² or heavier

Colors: Color Print on White

Printer: HP Color LaserJet 8550-PS

Finish: None

Adhesive: None

Special Notes: Illustrations are Ref Only ** Not to Scale ** (Page 1 of 2)



Application Note

How the LP80 Measures Leaf Area Index

Leaf area index (LAI) is just a single number—a statistical snapshot of a canopy taken at one particular time. But that one number can lead to significant insight, because it can be used to model and understand key canopy processes, including radiation interception, energy conversion, momentum, gas exchange, precipitation interception, and evapotranspiration.

Leaf area index is defined as the one-sided green leaf area of a canopy or plant community per unit ground area. It can therefore be found by harvesting and measuring the area of every leaf in a canopy covering one unit area of ground. In 1971, Anderson developed a less destructive method for finding LAI. Using hemispherical photographs looking upwards, he estimated the fraction of light that penetrated the canopy and applied a predictive mathematical model to approximate leaf area index.

Evaluating "fish eye" canopy pictures was tedious work. An assistant would usually lay a grid over each picture and count what fraction of the squares were light. One lab tech recalls, "After too many hours looking at those pictures, I used to dream in checkers." The "checkers" evaluation allowed investigators to find the probability that a random beam of light would penetrate that particular section of canopy.

The mathematical model that converts this fraction of light into an estimate of leaf area index is relatively simple. To understand how it works, picture holding a leaf with an area of ten square centimeters horizontally over a large white square. It would cast a shadow of ten square centimeters. Then, randomly place an identically sized leaf over the square. In all probability, the shadow cast would now be twenty square centimeters, although there is a small chance that the leaves might overlap. When a third leaf is added, the probability of overlap increases, and as more and more leaves are randomly placed, eventually the white square will be completely shaded, and though leaf area will increase as leaves are added, the shaded area will remain constant because all light has been intercepted.

LAI Conversion

Getting a value for leaf area index is often just a point along the way. If you plan to use LAI to model environmental interactions of the canopy, measuring photosynthetically active radiation (PAR) may be a more direct route. That's because many of those models are using LAI to predict PAR in the first place. It's possible to go back the other way—to use PAR to estimate LAI. But why do that if PAR is the number you really want? You may want to evaluate whether LAI is the most useful parameter in your particular application. It is sometimes more straightforward, and usually more accurate, to simply measure intercepted PAR and use that data directly in an appropriate model.

The equation describing this phenomenon (see Solving the Equation below for its mathematical derivation) is

$$t = \exp(-K \cdot LAI)$$

t is the probability that a ray will penetrate the canopy, LAI is the leaf area index of the canopy, and K is the extinction coefficient of the canopy. If you measure photosynthetically active radiation both above and below a canopy on a bright sunny day, the ratio of the two (PAR below to PAR above) is approximately equal to t. If you know K, you can find leaf area index (LAI) by inverting the equation:

$$LAI = -\ln t / K$$

The LP80 basically solves this equation to find leaf area index. But there are a couple of complicating factors. In constructing the model, we assumed that the leaves in our artificial canopy were horizontal and black, and that all radiation came directly from the sun. In reality, the angle of the sun changes over the course of the day, and real canopies have quite complex architecture. Also, some radiation is scattered both from leaves in the canopy and from the sky. A full model for finding the leaf area index