



METER  
ENVIRONMENT

## HOW TO ASSESS MAXIMUM POTENTIAL BIOMASS PRODUCTION

Dr. Gaylon S. Campbell

The conversion of light energy and atmospheric carbon dioxide to plant biomass is fundamentally important to both agricultural and natural ecosystems. The detailed biophysical and biochemical processes by which this occurs are well understood. At a less-detailed level, however, it is often useful to have a simple model that can be used to understand and analyze parts of an ecosystem. Such a model has been provided by Monteith (1977). He observed that when biomass accumulation by a plant community is plotted as a function of the accumulated solar radiation intercepted by the community, the result is a straight line. Figure 1 shows Monteith's results.




Figure 1. Total dry matter produced by a crop as a function of total intercepted radiation (from Monteith, 1977)

## QUANTIFY THE EFFECT OF PLANT STRESS

The mathematical model suggested by Figure 1 is

$$A = e f S_t$$

Equation 1

where  $A$  is biomass accumulation,  $S_t$  is the total solar radiation incident on the canopy,  $f$  is the fraction of incident radiation intercepted by the canopy, and  $e$  is the conversion efficiency for the canopy. For this calculation,  $A$  typically has units of  $\text{kg m}^{-2} \text{day}^{-1}$ . When  $S_t$  is the total solar radiation in  $\text{MJ/day}$ ,  $e$  has a value of around  $0.015 \text{ kg/MJ}$  for C3 crop species.

A number of experiments have shown that  $e$  is very conservative in situations where water, nutrients, and temperature do not limit plant growth. Equation 1 is therefore useful for predicting maximum productivity. When stresses limit growth, it is often possible to quantify their effect either in terms of a reduction in conversion efficiency,  $e$ , or a decrease in interception,  $f$ . This allows experiments carried out under different conditions of light availability to be compared or normalized.

The [ACCUPAR LP-80](#) can be used to measure  $f$ , the fraction of incident radiation intercepted by the canopy.

## REFERENCE

Monteith, James L., and C. J. Moss. "Climate and the efficiency of crop production in Britain." *Philosophical Transactions of the Royal Society of London B: Biological Sciences* 281, no. 980 (1977): 277-294. [Article link](#)

## QUESTIONS?

Explore questions and ideas with a canopy expert. METER scientists have over 100 years combined experience measuring the soil-plant-atmosphere continuum. Learn more about canopy measurement in the video [here](#). Dr. Steve Garrity discusses Leaf Area Index (LAI). Topics covered include the theory behind the measurement, direct

and indirect methods, variability among those methods, things to consider when choosing a method, and applications of [LAI](#).

## GET THE COMPLETE PICTURE

Learn more about measuring canopy. Get everything you need to know about measuring leaf area index, all in one place.

[Download the “Researcher’s complete guide to leaf area index \(LAI\)”](#)

REQUEST A QUOTE

CONTACT US